

Patientinformation

Losartan	Cozaar/Comp, Klomentan/Comp, Losarstad/Comp, Losartan, Losatrix, Losartan/Hydrochlorothiazide, Losartan/hydroklortiazid, Marozid
Eprosartan	Teveten/Comp
Valsartan	Diovan/Comp, Valsartan, Valsartan/hydrochlorothiazide, Valsartan/Hydroklortiazid Valsartore/Comp, Valtsu/Comp, Exforge, Entresto
Irbesartan	Aprovel, CoAprovel, Ifirmasta, Ifirmacombi, Irbesartan, Irbesartan/hydrochlorothiazid, Karvea
Kandesartan	Atacand/Plus, Candemox Comp, Candesarstad/Comp, Candesartan, Candesartan/Hydrochlorothiazid, Candexetil/Comp, Kairasec, Kandrozid,
Telmisartan	Micardis/Plus, Lesamor, Telmisartan/Hydroklortiazid, Tolura, Tolucombi
Diabetes medicines	
Metformin	Metformin, Comptact, Eucreas, Glucophage, Janumet, Jentaduetto, Komboglyze, Synjardy, Vokanamet, Xigduo
Empagliflozin	Jardiance, Synjardy, Glyxambi
Kanagliflozin	Invokana, Vokanamet
Dapagliflozin	Forxiga, Xigduo, Qtern
Ertugliflozin	Steglatro, Segluromet, Steglujan
Other medicines	

Painkillers in gel or spray form can be taken. Tablets and liquid medicines containing paracetamol (Alvedon, Pamol, Panodil, Paracetamol, Paracut, Pinex, Quramol, Theremin) can also be taken.

Never exceed the recommended dose!

Produced by the Elderly and Medication Therapy Group
(Terapigrupp Äldre och Läkemedel), August 2018, Version 2

**If you are taking
MEDICINES
that carry a risk of serious side effects in
conjunction with acute dehydration caused by**

- *Diarrhoea*
- *Vomiting*
- *High fever (over 38°C)*
- *Heatwave*
- *Acute illness resulting in difficulty eating or drinking*

Certain medicines that are used to treat heart conditions, high blood pressure, pain and diabetes could have serious side effects if they are taken when your body's fluid balance is disrupted.

Remember:

There is a risk of serious side effects from these medicines in conjunction with conditions that cause dehydration, such as vomiting, diarrhoea, fever, strong heat, or a disorder that renders you unable to eat or drink. The risk of serious side effects increases for elderly persons or individuals with reduced kidney function.

What action should be taken in the event of dehydration?

If you are taking any of the named medicines, you ought to discontinue treatment temporarily if you are dehydrated. If the problem is only temporary, treatment can recommence once it has ceased.

Contact your healthcare provider if you are in any doubt!

.....

If you are taking:

Painkillers		
Diklofenak	Diclofenak/T, Dicuno, Diklofenak/T, Eeze, Eezeneo, Ignorin, Voltaren/T, Arthrotec/forte	
Piroxikam	Brexidol	
Tenoxikam	Alganex	
Meloxikam	Meloxicam	
Ibuprofen	Brufen/retard, Ibumax, Ibumetin, Ibuprofen, Ibuzin, Ifenin, Ipren, Nurofen Apelsin	
Naproxen	Alpoxen, Naprocur, Naprosyn Entero, Naproxen, Pronaxen, Vimovo	
Ketoprofen	Orudis/Retard	
Dexibuprofen	Tradil	
Dexketoprofen	Enantyum, Kettesse	
Celecoxib	Celebra, Celecoxib	
Etoricoxib	Arcoxia, Turox	
Nabumeton	Relifex	
Diuretics		
Bendroflumetiazid	Bendroflumetiazid, Salures, Centyl K/mite	
Hydroklortiazid	Esidrex, Hydrochlorothiazide, Hydroklortiazid	
Clortalidon	Hygropax	
Metolazon	Metolazon	
Furosemid	Furix/Retard, Furosemid, Impugan, Lasix Retard	
Bumetanid	Burinex	
Torasemid	Torasemid, Torem	
Spirololakton	Spirolactone, Spirolakton	
Eplerenon	Eplerenon, Inspra	
Amilorid	Amilorid	
Medicines for heart conditions and high blood pressure		
Digoxin	Digoxin, Lanoxin	
Kaptopril	Captopril	
Enalapril	Enalapril, Renitec, Enalapril Comp, Enalapril/Hydrochlorothiazide, Linatil Comp, Linatil Comp Mite, Renitec Comp	
Lisinopril	Lisinopril, Zestril, Lisinopril/Hydrochlorotiazid	
Perindopril	Coversyl Novum	
Ramipril	Ramipril, Triatec, Ramipril/Hydroklorotiazid	
Fosinopril	Fosinopril	
Kinapril	Accupro Comp	